


Use the following information to answer the first question.




1. The activation energy for the forward, catalyzed reaction is

- (A) II minus I
- B. III minus I
- C. IV minus II
- D. IV minus III

Use the following information to answer the next question.

Incomplete combustion in motor vehicles may lead to the formation of carbon monoxide gas, which is a health hazard in high concentrations. Carbon monoxide gas is converted to carbon dioxide gas in a catalytic converter before being emitted from the motor vehicle. This conversion is represented by the equation below.



2. The addition of a catalyst to the reaction represented by the equation above would i the energy transferred during the reaction and would ii the value of the equilibrium constant.

The statement above is completed by the information in row

Row	i	ii
A.	increase	increase
B.	increase	not change
C.	not change	increase
(D)	not change	not change

Use the following information to answer the next question.

Honey has a high concentration of fructose, $\text{C}_6\text{H}_{12}\text{O}_6\text{(s)}$. Fructose has the same molecular formula as glucose but a different structural formula.

3. If 1.50 mmol of fructose is burned in a calorimeter that contains 250.0 g of water and the temperature increases by 3.85 °C, then the molar enthalpy of combustion of fructose is

- A. -6.05×10^{-3} kJ/mol
- B. -9.68×10^{-2} kJ/mol
- C. -4.03 kJ/mol
- (D) -2.69×10^3 kJ/mol

$$\Delta_c H = Q$$

$$n \Delta_c H_m = m c \Delta T$$

$$\Delta_c H_m = \frac{m c \Delta T}{n}$$

$$= \frac{(250\text{g})(4.19\text{J/g}\cdot^\circ\text{C})(3.85^\circ\text{C})}{n}$$


$$= 2688.5833\text{ J/mol}$$

$$0.00150$$

$$= 2688.5\text{ kJ/mol}$$

Use the following information to answer the next question.

A technician performed an experiment to determine the molar enthalpy of combustion of propane in a soldering torch, as represented in the diagram below.



4. If the experimental value of the molar enthalpy of combustion of propane in the technician's calorimetry experiment is significantly different from the theoretical value, then the technician could reduce the discrepancy in the data by

- A. using a glass beaker to hold the water
- B. creating an enclosing shield around the apparatus
- C. raising the aluminum calorimeter above the flame
- D. decreasing the mass of water in the aluminum calorimeter

5. During a combustion reaction, energy is i the surroundings because the products have ii potential energy than the reactants.

The statement above is completed by the information in row

Row	i	ii
<input checked="" type="radio"/> A.	released to	lower
B.	released to	higher
C.	absorbed from	lower
D.	absorbed from	higher

Use the following information to answer the next question.

Chemicals

- | | | | |
|---|-----------|---|--------------------|
| 1 | $O_2(g)$ | 4 | $H_2O(l)$ |
| 2 | $CO(g)$ | 5 | $H_2O(g)$ |
| 3 | $CO_2(g)$ | 6 | $C_6H_{12}O_6(aq)$ |


Numerical Response

1. Match the chemicals numbered above with the statements given below.

The reactants of photosynthesis are: 3 and 4
Record in the first column Record in the second column

The products of complete hydrocarbon combustion in an open system are: 3 and 5
Record in the third column Record in the fourth column

Use the following information to answer the next question.



6. The reaction represented in the diagram above is I, and if the energy was included as a term in the balanced equation, it would be a ii.

The statement above is completed by the information in row

Row	<i>i</i>	<i>ii</i>
A.	exothermic	reactant
B.	exothermic	product
<input checked="" type="radio"/> C.	endothermic	reactant
D.	endothermic	product


Use the following information to answer the next two questions.

The bombardier beetle can release a chemical solution when threatened. Glands in the beetle produce hydrogen peroxide and hydroquinone, $C_6H_4(OH)_2(aq)$, which are combined to produce the reaction represented by the overall equation below.



The equations listed below represent reactions that are related to the production of the chemical solution.

Equations



7. The enthalpy change for the overall equation is

- A. +83.0 kJ
 B. -202.6 kJ
 C. -299.4 kJ
 D. -585.0 kJ

Use the following additional information to answer the next question.





8. Which of the following substances would oxidize $C_6H_4(OH)_2(aq)$?

- A. $Ag^+(aq)$
 B. $Cl^{3+}(aq)$
 C. $Ag(s)$
 D. $Cl(s)$

Use the following information to answer the next two questions.

Nitrogen can react with oxygen to form a variety of oxides as represented by the following equations.



Numerical Response

2. The oxidation number of nitrogen in

NO(g) is 2 (Record in the first column)

NO₂(g) is 4 (Record in the second column)

N₂O(g) is 1 (Record in the third column)

N₂O₄(g) is 4 (Record in the fourth column)

Use the following additional information to answer the next question.




Numerical Response

3. The nitrogen oxides, listed in order of increasing enthalpy of formation, are 4, 2, 3, and 1.

Use the following information to answer the next three questions.

The energy from a car battery is generated as represented by the equation below.



$$\Delta H = -315.9 \text{ kJ}$$

9. If 15.0 g of Pb(s) reacts in a car battery, the amount of energy released is

A. 4.74 MJ

B. 4.36 MJ

C. 22.9 kJ

D. 21.1 kJ

$$\Delta_r H_{\text{m}} = n \Delta_r H_{\text{m}}$$

$$= \left(\frac{15.0}{207.20} \text{ mol} \right) (-315.9 \text{ kJ/mol}) = -22.869 \text{ kJ} \approx -22.87 \text{ kJ}$$

$$\Delta_r H_{\text{m}} = \frac{-315.9 \text{ kJ}}{\text{Pb}} \times 1 \text{ mol} = -315.9 \text{ kJ/mol}$$

10. During the operation of a car battery, which of the following observations can be made?

A. The amount of Pb(s) increases as PbO₂(s) is reduced.

B. The amount of PbO₂(s) increases as Pb(s) is reduced.

C. The amount of PbO₂(s) decreases as Pb(s) is oxidized.

D. The amount of Pb(s) decreases as PbO₂(s) is oxidized.

Use the following additional information to answer the next question.

Every car battery is given a CCA (cold cranking amps) rating. A CCA rating of 600 means that the battery is capable of generating 600 A of current for a 30.0 s period at 0 °C.

11. Which of the following values indicates how many coulombs a battery with a CCA rating of 600 produces during 30.0 s of operation?

A. 20.0 C

B. 600 C

C. 1.80 × 10⁴ C

D. 1.74 × 10⁹ C


$$Q = I t$$

$$= (600 \text{ A})(30 \text{ s})$$

Use the following information to answer the next two questions.

The electrolysis of aluminum oxide in an electrolytic cell occurs at high temperatures so that the compound is molten.

12. Which of the following equations represents the reduction half-reaction when molten aluminum oxide undergoes electrolysis?




13. During the production of aluminum metal in the electrolytic cell, anions travel toward the *i* and electrons travel through the *ii* .

The statement above is completed by the information in row


Row	<i>i</i>	<i>ii</i>
A.	cathode	electrolyte to the anode
B.	cathode	wire to the cathode
C.	anode	electrolyte to the anode
D.	anode	wire to the cathode

Use the following information to answer the next question.

A student constructs the following standard electrochemical cell in a laboratory at 25.0 °C.



14. If the standard lead reduction half-reaction had been chosen as the reference half-reaction instead of the hydrogen reduction half-reaction, then the electrical potential for this cell would be





$$E_{cell} = E_{cath} - E_{anode}$$

$$= +0.46 V$$

Use the following information to answer the next two questions.

In an experiment to study the reactivity of Pt(s), Rh(s), Sm(s), and Te(s), a student observed the reactions represented by the equations below.



15. Which of the following substances is the strongest reducing agent?

- A. Pt(s)
- B. Rh(s)
- C. Sm(s)
- D. Te(s)

16. Which of the following equations represents a spontaneous reaction?



Use the following information to answer the next two questions.

A student placed a large piece of zinc into a beaker of hydrochloric acid and collected all of the gas produced. Indicators were also added to monitor the change in pH.

17. Which of the following rows gives the composition of the bubbles and the process through which they were formed?




Row	Composition	Process of Formation
A.	$\text{Cl}_2(\text{g})$	oxidation of chloride ions
<input checked="" type="radio"/> B.	$\text{H}_2(\text{g})$	reduction of hydrogen ions
C.	$\text{H}_2(\text{g})$	reduction of water
D.	$\text{O}_2(\text{g})$	oxidation of water

18. If a student were to build a voltaic cell using solid zinc and hydrochloric acid, which of the following equipment would also be needed?

- A. An inert electrode for the cathode and a salt bridge
- B. An inert electrode for the cathode and a power source
- C. An inert electrode for the anode and a salt bridge
- D. An inert electrode for the anode and a power source





Use the following information to answer the next question.

Fireworks usually contain a mixture of explosives and other chemicals. Some of the reactions that occur in a fireworks display are represented by the equations below.



19. The equations above that represent a reaction in which the metal is being oxidized are

- A. I and IV only
- B. II and III only
- C. I, III, and IV
- D. II, III, and V



Use the following information to answer the next question.

Ammonium nitrate, used to make gunpowder and fireworks, was extracted from animal manure in ancient China. During the explosion of gunpowder or fireworks, the ammonium nitrate reacts violently, as represented by the equation below.


Explosion of Ammonium Nitrate



20. During the explosion of ammonium nitrate, hydrogen
- A. is oxidized
 - B. loses electrons
 - C. is the oxidizing agent
 - B** has no change in oxidation number

Use the following information to answer the next question.


Methanolic acid, in the presence of a catalyst, can be used to produce electricity in a fuel cell, as represented by the following diagram.



21. The equation that represents the half-reaction that occurs at the cathode of the fuel cell is
- A** $\text{O}_2(g) + 4\text{H}^+(aq) + 4e^- \rightarrow 2\text{H}_2\text{O}(l)$
 - ~~B~~ $2\text{H}_2\text{O}(l) \rightarrow \text{O}_2(g) + 4\text{H}^+(aq) + 4e^-$
 - ~~C~~ $\text{HCOOH}(l) \rightarrow \text{CO}_2(g) + 2\text{H}^+(aq) + 2e^-$
 - D. $\text{CO}_2(g) + 2\text{H}^+(aq) + 2e^- \rightarrow \text{HCOOH}(l)$

Use the following information to answer the next question.

The equation below represents the $\text{AlF}_6^{3-}(aq)$ reduction half-reaction.



One half-cell in an electrochemical cell contains Al(s) in a $\text{F}^-(aq)$ solution. The other half-cell contains Pb(s) in a $\text{Pb}^{2+}(aq)$ solution. A spontaneous reaction occurs, producing $\text{AlF}_6^{3-}(aq)$ and Pb(s).


Numerical Response

4. The net cell potential for this electrochemical cell is +/- 1.94 V.

$$E_{\text{cell}} = E_{\text{cath}} - E_{\text{anode}}$$

$$= (-0.13\text{V}) - (-2.07\text{V})$$
$$= 1.94\text{V}$$

Use the following information to answer the next question.



22. If the electrochemical cell in the diagram above produces a flow of electrons in the direction indicated, then $M(s)$ and $M^{2+}(aq)$ could be

- (A) $Fe(s)$ and $Fe^{2+}(aq)$
 B. $Pb(s)$ and $Pb^{2+}(aq)$
 C. $Ni(s)$ and $Ni^{2+}(aq)$
 D. $Cu(s)$ and $Cu^{2+}(aq)$

$M(s)$ has to be below
 $Cd(aq)$ in order for a spontaneous reaction
 to occur

Use the following information to answer the next two questions.

Statements About Electrochemical Cells

- I The reaction is spontaneous.
 (II) The reaction is nonspontaneous.
 Anions migrate to the anode.
 (III) Cations migrate to the anode.
 IV Electrons are gained at the anode.
 V Electrons are gained at the cathode.
 (VI) Electrons are gained at the cathode.

23. The statements above that correctly describe an electrolytic cell are

- A. I, III, and V
 B. I, IV, and VI
 (C) II, III, and VI
 D. II, IV, and V

24. The statements above that correctly describe both an electrolytic cell and a voltaic cell are

- A. I and III
 (B) III and VI
 C. IV and V
 D. IV and VI

Use the following information to answer the next question.

Iron metal reacts with hydrochloric acid slowly. The equation for this reaction is




25. In this reaction, the reducing agent is

- A. $FeCl_2(aq)$
 B. $HCl(aq)$
 C. $H_2(g)$
 (D) $Fe(s)$

→ gets oxidized

Use the following information to answer the next question.

A student drew the structural diagram shown below:



26. The IUPAC name for the structural diagram the student drew is 1- i - 2 - ii .

The statement above is completed by the information in row

Row	i	ii
A.	methyl	ethylcane
B.	methyl	ethylcyclobutane
C.	ethyl	methylbutane
D.	ethyl	methylcyclobutane

Use the following information to answer the next question.

Organic Compounds


- | | | | |
|---|------------------------|---|--------------------|
| 1 | 2-methylcyclobut-1-ene | 4 | 5-methylhept-3-yne |
| 2 | 1,2-dibromohexane | 5 | cycloheptane |
| 3 | 2,2-dimethylpentane | 6 | pentan-1-ol |

Numerical Response

5. The organic compound numbered above that is an alkene is 1 (Record in the first column)
 is an alcohol is 6 (Record in the second column)
 contains a triple bond is 4 (Record in the third column)
 is cyclic and saturated is 5 (Record in the fourth column)

Use the following information to answer the next question.

Organic Compounds



Numerical Response

6. Match four of the organic compounds numbered above with their classifications below.
- | | | |
|-------------------------|--------------|-------------------------------|
| Alkyne | <u> 4 </u> | (Record in the first column) |
| Alcohol | <u> 6 </u> | (Record in the second column) |
| Aromatic | <u> 1 </u> | (Record in the third column) |
| Unsaturated hydrocarbon | <u> 3 </u> | (Record in the fourth column) |

Use the following information to answer the next question.

Reaction Equation



Names and Terms

- | | |
|---------------------|------------------|
| 1 Methane | 6 Ester |
| 2 Methanol | 7 Polymer |
| 3 Ethanoate | 8 Esterification |
| 4 Methanoic acid | 9 Polymerization |
| 5 Methyl methanoate | |


Numerical Response

7. Match a name or a term from the list above with each descriptor given below.

- Name of reactant I 4 (Record in the first column)
 Name of product II 5 (Record in the second column)
 Type of reaction 8 (Record in the third column)
 Classification of product II 6 (Record in the fourth column)

Use the following information to answer the next question.

The following are structural diagrams for four organic compounds with common industrial uses.



Numerical Response

8. Match each of the structural diagrams above with its classification below.

- Aromatic 1 (Record in the first column)
 Carboxylic acid 3 (Record in the second column)
 Unsaturated and aliphatic 4 (Record in the third column)
 Halogenated hydrocarbon 2 (Record in the fourth column)

Use the following information to answer the next two questions.

Hexane and hex-1-ene are both colorless liquids. One method used to differentiate between hexane and hex-1-ene is to add a few drops of orange-coloured aqueous bromine to samples of each organic compound.

27. Hexane is I *hydration*, and hex-1-ene is II *hydrocarbon*.

The statement above is completed by the information in row

Row	I	II
A.	a saturated	a saturated
B.	a saturated	an unsaturated
C.	an unsaturated	a saturated
D.	an unsaturated	an unsaturated

28. When aqueous bromine is added to hexane and hex-1-ene in the presence of light, the hexane undergoes I *reaction* and the hex-1-ene undergoes II *reaction*.

The statement above is completed by the information in row

Row	I	II
A.	an addition	a substitution
B.	an addition	an addition
C.	a substitution	a substitution
D.	a substitution	an addition

Use the following information to answer the next question.

Carbon-Containing Compounds

1	$\text{CCl}_4(\text{l})$	5	$\text{CO}(\text{g})$
2	$\text{Fe}_2\text{Cl}_6(\text{s})$	6	$\text{C}_2\text{H}_4(\text{g})$
3	$\text{C}_2\text{H}_2(\text{g})$	7	$\text{NaCN}(\text{s})$
4	$\text{C}_2\text{H}_5\text{OH}(\text{l})$	8	$\text{MgCO}_3(\text{s})$

Numerical Response



9. The compounds numbered above that can be classified as organic are 1, 3, 4, and 6.

11/10
11/10
11/10

11/10
11/10
11/10

Use the following information to answer the next question.

Organic Compounds



29. An ester functional group is found in

- A. II and III only
 B. II, III, and IV
 C. III only
 D. V only

Use the following information to answer the next two questions.

The concentration of aqueous sodium hypochlorite, NaOCl(aq) , in laundry bleach can be determined by titrating a sample of laundry bleach with an iodide solution, as represented by the equation below.



$n = 1.405 \times 10^{-4} \text{ mol}$
 $V = 0.100 \text{ L}$
Numerical Response
 4.25 mL
 0.0473 mol/L
 $n = 2.01 \times 10^{-4} \text{ mol}$

10. If a student uses 4.25 mL of a 0.0473 mol/L $\text{I}^-(\text{aq})$ solution to titrate a 100.00 mL sample of laundry bleach, then the concentration of $\text{OCl}^-(\text{aq})$ in the laundry bleach is 1.00 mmol/L.


30. The K_b of $\text{OCl}^-(\text{aq})$ is 1, and $\text{OCl}^-(\text{aq})$ is a weaker base than H.

The statement above is completed by the information in row

Row	i	#
<input checked="" type="radio"/> A.	2.5×10^{-7}	$\text{PO}_4^{3-}(\text{aq})$
<input type="radio"/> B.	2.5×10^{-7}	$\text{CH}_3\text{COO}^-(\text{aq})$
<input type="radio"/> C.	4.0×10^{-5}	$\text{PO}_4^{3-}(\text{aq})$
<input type="radio"/> D.	4.0×10^{-8}	$\text{CH}_3\text{COO}^-(\text{aq})$

Use the following information to answer the next two questions.

Citric acid, $\text{H}_3\text{C}_6\text{H}_5\text{O}_7(\text{aq})$, is a weak, polyprotic acid that is found in fruits such as oranges and lemons. Citric acid reacts with water, as represented by the following Brønsted-Lowry equations.



$$\begin{aligned} K_a &= 1.35 \times 10^{-4} \\ K_b &= 5.88 \times 10^{-10} \\ K_b &= 2.5 \times 10^{-8} \end{aligned}$$

31. The amphiprotic species in the equations above are

- A. ~~$\text{H}_2\text{C}_6\text{H}_4\text{O}_7^-(\text{aq})$~~ , $\text{HC}_6\text{H}_3\text{O}_7^{2-}(\text{aq})$, and $\text{H}_2\text{O}(\text{l})$
 B. $\text{H}_2\text{C}_6\text{H}_4\text{O}_7^-(\text{aq})$, $\text{HC}_6\text{H}_3\text{O}_7^{2-}(\text{aq})$, and $\text{H}_2\text{O}(\text{l})$
 C. ~~$\text{H}_2\text{C}_6\text{H}_4\text{O}_7^-(\text{aq})$~~ and $\text{H}_2\text{C}_6\text{H}_4\text{O}_7^-(\text{aq})$
 D. $\text{HC}_6\text{H}_3\text{O}_7^{2-}(\text{aq})$ and $\text{C}_6\text{H}_5\text{O}_7^{3-}(\text{aq})$

32. Which of the following statements about K_a and K_b values applies to the equations above?

- A. The K_a of $\text{H}_2\text{C}_6\text{H}_4\text{O}_7^-(\text{aq})$ is less than the K_b of $\text{HC}_6\text{H}_3\text{O}_7^{2-}(\text{aq})$.
 B. The K_b of $\text{HC}_6\text{H}_3\text{O}_7^{2-}(\text{aq})$ is greater than the K_a of $\text{C}_6\text{H}_5\text{O}_7^{3-}(\text{aq})$.
 C. The K_a of $\text{H}_2\text{C}_6\text{H}_4\text{O}_7^-(\text{aq})$ is greater than the K_b of $\text{C}_6\text{H}_5\text{O}_7^{3-}(\text{aq})$.
 D. The K_b of $\text{H}_2\text{C}_6\text{H}_4\text{O}_7^-(\text{aq})$ is greater than the K_b of $\text{HC}_6\text{H}_3\text{O}_7^{2-}(\text{aq})$.

Use the following information to answer the next two questions.

Lactic acid, $\text{HC}_3\text{H}_5\text{O}_3(\text{aq})$, is produced in human muscle cells when not enough oxygen is supplied to the muscle during heavy physical activity. The equation below represents the Brønsted-Lowry reaction of lactic acid and water.



33. Which of the following rows identifies the Brønsted-Lowry acids and a conjugate acid-base pair in the equation above?

Row	Brønsted-Lowry Acids	Conjugate Acid-Base Pair
A.	$\text{HC}_3\text{H}_5\text{O}_3(\text{aq})$ and $\text{C}_3\text{H}_5\text{O}_3^-(\text{aq})$	$\text{H}_2\text{O}(\text{l})$ and $\text{H}_3\text{O}^+(\text{aq})$
B.	$\text{HC}_3\text{H}_5\text{O}_3(\text{aq})$ and $\text{C}_3\text{H}_5\text{O}_3^-(\text{aq})$	$\text{C}_3\text{H}_5\text{O}_3^-(\text{aq})$ and $\text{H}_3\text{O}^+(\text{aq})$
C.	$\text{HC}_3\text{H}_5\text{O}_3(\text{aq})$ and $\text{H}_3\text{O}^+(\text{aq})$	$\text{H}_2\text{O}(\text{l})$ and $\text{H}_3\text{O}^+(\text{aq})$
D.	$\text{HC}_3\text{H}_5\text{O}_3(\text{aq})$ and $\text{H}_3\text{O}^+(\text{aq})$	$\text{C}_3\text{H}_5\text{O}_3^-(\text{aq})$ and $\text{H}_3\text{O}^+(\text{aq})$

Use the following additional information to answer the next question.

A 100.0 mL sample of lactic acid has a pH of 3.38.

Numerical Response


34. The hydroxide ion concentration in this sample of lactic acid, expressed in scientific notation, is $a.b \times 10^{-cd}$ mol/L. The values of a , b , c , and d are 2, 9, 1 and 1.

$$\text{pOH} = 14 - \text{pH} < 10.62$$

$$[\text{OH}^-] = 10^{-\text{pOH}} = 2.39 \times 10^{-11}$$

34. If a 100.0 mL sample of 0.167 mol/L unknown acid has a pH of 2.32 at 25.0 °C, then the K_a is

- A. 2.9×10^{-2}
 B. 4.8×10^{-3}
 C. 1.4×10^{-4}
 D. 2.3×10^{-5}




$$K_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]}$$

Use the following information to answer the next question.

A technician placed an amount of the colourless gas dinitrogen tetraoxide into a flask. He closed the flask and allowed the reaction to reach equilibrium. The dinitrogen tetraoxide partially decomposed to form brown-coloured nitrogen dioxide gas. The data collected during the experiment were recorded below.

	$\text{N}_2\text{O}_4(\text{g})$	$\text{NO}_2(\text{g})$
Initial Concentration (mol/L)	0.700	0.000
Final Concentration (mol/L)	0.610	0.180

35. The balanced chemical equation and equilibrium constant for the partial decomposition of dinitrogen tetraoxide gas are



Use the following information to answer the next question.

When the system represented by the equation below is at equilibrium in a 2.00 L flask at 15.0 °C, the flask contains 1.15 mmol of $\text{H}_2(\text{g})$, 2.13 mmol of $\text{I}_2(\text{g})$, and 3.74 mmol of $\text{HI}(\text{g})$.



Numerical Response


12. At 15.0 °C, the equilibrium constant is 5.71

$$K_c = \frac{0.00187}{(5.75 \times 10^{-4})(0.001065)}$$

Use the following information to answer the next two questions.

In blood, the enzyme carbonic anhydrase catalyzes the formation of carbonic acid from aqueous carbon dioxide and water. Carbonic acid and hydrogen carbonate form an important buffer in the blood. Two reactions that occur in the blood are represented by the equations below.

Reactions in the Blood




36. If the concentration of $\text{CO}_2(\text{aq})$ in the blood increases, then the equilibria will shift to the i, and the concentration of $\text{HCO}_3^-(\text{aq})$ in the blood will ii.

The statement above is completed by the information in row

Row	i	ii
A.	left	increase
B.	left	decrease
C.	right	increase
D.	right	decrease

37. The equilibrium law expression for the reaction represented by equation II is.




Use the following information to answer the next question.

At 200 °C, the equilibrium system represented by the following equation and diagram was established.



Equilibrium System





38. In the equilibrium system represented in the diagram above, equilibrium was initially established at 1, and the stress applied to the system at 14.0 minutes was ii in temperature.

The statement above is completed by the information in row

Row	i	ii
A.	4.5 min	an increase
B.	4.5 min	a decrease
C.	14.0 min	an increase
D.	14.0 min	a decrease

39. Which of the following graphs represents the titration of a weak, polyprotic base with a strong, monoprotic acid?



40. Which of the following systems could be at equilibrium?
- A.** A closed bottle of carbonated water
 - B. A block of ice in a glass of water
 - C. Water boiling in a kettle
 - D. A glass of pop

Use the following information to answer the next question.

Equations	K_c at 25 °C
1 $\text{H}_2(\text{g}) + \text{Br}_2(\text{g}) \rightleftharpoons 2 \text{HBr}(\text{g})$	5.0×10^{-18}
2 $\text{H}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightleftharpoons 2 \text{HCl}(\text{g})$	2.5×10^{15}
3 $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2 \text{NO}(\text{g})$	2.0×10^{-11}
4 $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2 \text{HI}(\text{g})$	2.5×10^{-1}

Numerical Response

13. When the equations numbered above are ordered from the reaction that produces the most products to the reaction that produces the least products, the order is

Most 4, 1, 3, and 2 Least

Use the following information to answer the next question.

Weak Acids
1 $\text{HF}(\text{aq})$
2 $\text{H}_2\text{S}(\text{aq})$
3 $\text{HOCl}(\text{aq})$
4 $\text{H}_2\text{SO}_3(\text{aq})$

Numerical Response

14. When the weak acids numbered above are ordered from the acid with the strongest conjugate base to the acid with the weakest conjugate base, the order is

Strongest 3, 2, 1, and 4 Weakest

Use the following information to answer the next question.

Pairs of Solutions
I $\text{HCl}(\text{aq})$ and $\text{NaOH}(\text{aq})$
II $\text{HClO}_4(\text{aq})$ and $\text{KClO}_4(\text{aq})$
III $\text{H}_2\text{SO}_4(\text{aq})$ and $\text{LiHSO}_4(\text{aq})$
IV $\text{H}_3\text{PO}_4(\text{aq})$ and $\text{NaH}_2\text{PO}_4(\text{aq})$

41. If each pair of solutions listed above is mixed together in equal amounts, then the pair of solutions that would act as a buffer is

- A. I
B. II
C. III
D. IV