


Which of the following changes is **not** an example of oxidation?

- A. Corrosion of metals
- B. Plating of metals
- C. Rusting of iron
- D. Reaction at the anode of a battery

An equation that represents a redox reaction is

- A.  $\text{NaOH}_{(aq)} + \text{HCl}_{(aq)} \rightarrow \text{NaCl}_{(aq)} + \text{H}_2\text{O}_{(l)}$
- B.  $\text{AgNO}_3_{(aq)} + \text{KI}_{(aq)} \rightarrow \text{AgI}_{(s)} + \text{KNO}_3_{(aq)}$
- C.  $\text{Mg}(\text{OH})_2_{(s)} + \text{H}_2\text{SO}_4_{(aq)} \rightarrow \text{MgSO}_4_{(aq)} + 2\text{H}_2\text{O}_{(l)}$
- D.  $\text{Cu}_{(s)} + 4\text{HNO}_3_{(aq)} \rightarrow \text{Cu}(\text{NO}_3)_2_{(aq)} + 2\text{NO}_2_{(g)} + 2\text{H}_2\text{O}_{(l)}$

Use the following equation to answer the next question.



### Numerical Response

When balanced in terms of lowest whole numbers, the coefficients for this equation are, respectively, \_\_\_\_, \_\_\_\_, \_\_\_\_, and \_\_\_\_.  
(Record all four digits on the answer sheet.)

A spontaneous reaction would occur between a 1.0 mol/L  $\text{Fe}^{3+}_{(aq)}$  solution and

- A.  $\text{I}_2_{(s)}$
- B.  $\text{Zn}_{(s)}$
- C.  $\text{Hg}_{(l)}$
- D. 1.0 mol/L  $\text{Fe}^{2+}_{(aq)}$

### Numerical Response

8. The volume of 0.160 mol/L  $\text{K}_2\text{Cr}_2\text{O}_7_{(aq)}$  required to completely react with 10.0 mL of acidic 0.881 mol/L  $\text{H}_2\text{O}_2_{(aq)}$  is \_\_\_\_\_ mL.

(Record your answer to three digits on the answer sheet.)

### Numerical Response

9. The oxidation numbers of carbon in  $\text{HCOOH}_{(aq)}$ ,  $\text{C}_6\text{H}_{12}\text{O}_6_{(s)}$ ,  $\text{CO}_2_{(g)}$ , and  $\text{CHCl}_3_{(g)}$ , respectively, are \_\_\_\_\_.

(Record all four digits on the answer sheet.)

The reaction  $2 \text{H}_2\text{O}_{(l)} \rightarrow 2 \text{H}_{2(g)} + \text{O}_{2(g)}$  is an example of an

- A. exothermic redox reaction
- B. endothermic redox reaction
- C. exothermic reaction that absorbs energy
- D. endothermic reaction that releases energy

In which change are electrons gained?

- A.  $\text{Ca}^{2+}_{(aq)}$  to  $\text{Ca}_{(s)}$
- B.  $2 \text{Cl}^{-}_{(aq)}$  to  $\text{Cl}_{2(g)}$
- C.  $\text{Fe}^{2+}_{(aq)}$  to  $\text{Fe}^{3+}_{(aq)}$
- D.  $\text{NaCl}_{(s)}$  to  $\text{Na}^{+}_{(aq)}$  and  $\text{Cl}^{-}_{(aq)}$

Which ion could **not** act as both an oxidizing agent and a reducing agent?

- A.  $\text{Cu}^{2+}_{(aq)}$
- B.  $\text{Sn}^{2+}_{(aq)}$
- C.  $\text{Fe}^{2+}_{(aq)}$
- D.  $\text{Cr}^{2+}_{(aq)}$

Bacteria in our mouths and digestive systems convert sodium nitrate and other nitrate salts into nitrites, as indicated by the incomplete and unbalanced half-reaction  $\text{NaNO}_{3(aq)} \rightarrow \text{NaNO}_{2(aq)}$ . In this half-reaction,

- A. the oxidation number for sodium changes from  $-9$  to  $-7$
- B. the oxidation number of nitrogen increases
- C. the oxidation number of oxygen increases
- D. reduction occurs

Two reagents that will oxidize  $\text{Pb}_{(s)}$  to  $\text{Pb}^{2+}_{(aq)}$  but that will **not** oxidize  $\text{I}^{-}_{(aq)}$  to  $\text{I}_{2(s)}$  are

- A.  $\text{F}_{2(g)}$  and  $\text{Fe}^{3+}_{(aq)}$  ✓
- B.  $\text{Fe}^{3+}_{(aq)}$  and  $\text{Br}_{2(l)}$
- C.  $\text{Cd}^{2+}_{(aq)}$  and  $\text{Ag}^{+}_{(aq)}$
- D.  $\text{Cu}^{2+}_{(aq)}$  and  $\text{Sn}^{4+}_{(aq)}$

Metals  $W_{(s)}$ ,  $X_{(s)}$ ,  $Y_{(s)}$ , and  $Z_{(s)}$  were placed in solutions of each of their respective ionic salts. The results are summarized in the data table. A check mark indicates that a reaction occurred.

	$X^+_{(aq)}$	$Y^{2+}_{(aq)}$	$Z^{3+}_{(aq)}$	$W^+_{(aq)}$
$X_{(s)}$	—	no reaction	✓	✓
$Y_{(s)}$	✓	—	✓	✓
$Z_{(s)}$	no reaction	no reaction	—	✓
$W_{(s)}$	no reaction	no reaction	no reaction	—

According to the results, the strongest reducing agent is

- A.  $Y^{2+}_{(aq)}$
- B.  $W_{(s)}$
- C.  $Y_{(s)}$
- D.  $W^+_{(aq)}$

For the standard reference half-cell, the oxidation half-reaction and  $E^\circ$  are

- A.  $H_{2(g)} \rightarrow 2H^+_{(aq)} + 2e^-$   $E^\circ = 0.00\text{ V}$
- B.  $2H^+_{(aq)} + 2e^- \rightarrow H_{2(g)}$   $E^\circ = 0.00\text{ V}$
- C.  $2H_2O_{(l)} + 2e^- \rightarrow H_{2(g)} + 2OH^-_{(aq)}$   $E^\circ = -0.83\text{ V}$
- D.  $H_{2(g)} + 2OH^-_{(aq)} \rightarrow 2e^- + 2H_2O_{(l)}$   $E^\circ = +0.83\text{ V}$


Electrolytic cells are used commercially in

- A. cameras
- B. fuel cells
- C. flashlights
- D. metal plating

Which of the following aqueous ions can either gain or lose electrons in a redox reaction?

- A.  $Sn^{2+}_{(aq)}$
- B.  $Cl^-_{(aq)}$
- C.  $Ca^{2+}_{(aq)}$
- D.  $S^{2-}_{(aq)}$

If the lithium reduction half-reaction,  $Li^+_{(aq)} + e^- \rightarrow Li_{(s)}$ , had been assigned an  $E^\circ$  value of 0.00 V, the predicted  $E^\circ_{net}$  value for the reaction




- A. +3.38 V
- B. -2.28 V
- C. -0.42 V
- D. -1.10 V

The equation representing a spontaneous reaction at standard conditions is


- A.  $Co^{2+}_{(aq)} + 2Fe^{2+}_{(aq)} \rightarrow Co_{(s)} + 2Fe^{3+}_{(aq)}$
- B.  $Sn^{4+}_{(aq)} + 2Br^-_{(aq)} \rightarrow Sn^{2+}_{(aq)} + Br_{2(l)}$
- C.  $2I^-_{(aq)} + Cl_{2(g)} \rightarrow I_{2(s)} + 2Cl^-_{(aq)}$
- D.  $Pb_{(s)} + Fe^{2+}_{(aq)} \rightarrow Pb^{2+}_{(aq)} + Fe_{(s)}$

Use the following information to answer the next 5 questions

Silver oxide cells are efficient but expensive because they contain silver. The diagram illustrates the construction of a silver oxide cell.



The half reactions are



The anode of the cell is

- A.  $\text{Ag}_{(s)}$
- B.  $\text{Zn}_{(s)}$
- C.  $\text{Ag}_2\text{O}_{(s)}$
- D.  $\text{Zn}(\text{OH})_{2(s)}$

Using lowest whole number coefficients, the coefficient for  $\text{H}_2\text{O}_{(l)}$  in the balanced oxidation-reduction reaction that occurs during discharging of the cell is

- A. 1
- B. 2
- C. 3
- D. 4

As the cell operates, the species oxidized is

- A.  $\text{Ag}_{(s)}$
- B.  $\text{Zn}_{(s)}$
- C.  $\text{Ag}_2\text{O}_{(s)}$
- D.  $\text{Zn}(\text{OH})_{2(s)}$


As the cell operates, the

- A.  $[\text{OH}^-_{(aq)}]$  increases
- B. mass of  $\text{Zn}_{(s)}$  increases
- C. mass of  $\text{Ag}_2\text{O}_{(s)}$  decreases
- D. mass of  $\text{Zn}(\text{OH})_{2(s)}$  decreases

### Numerical Response

The voltage generated by the silver oxide cell is \_\_\_\_\_ V.

Use the following to answer the next question




**Numerical Response**

What reagents are required in order for the cell to produce a voltage of 1.25 V?

- Electrode I \_\_\_\_\_ (Record in first column)  
 Solution II \_\_\_\_\_ (Record in second column)  
 Electrode III \_\_\_\_\_ (Record in third column)  
 Solution IV \_\_\_\_\_ (Record in fourth column)

Use the following information to answer the next two questions.



The cell shown in the diagram is


- A. electrolytic
- B. voltaic
- C. galvanic
- D. acid-base

Products I, II, and III from this cell, respectively, are

- A.  $\text{Cl}_{2(g)}$ ,  $\text{H}_{2(g)}$ , and  $\text{HCl}_{(aq)}$
- B.  $\text{H}_{2(g)}$ ,  $\text{Cl}_{2(g)}$ , and  $\text{NaOH}_{(aq)}$
- C.  $\text{HCl}_{(g)}$ ,  $\text{Cl}_{2(g)}$ , and  $\text{NaOH}_{(aq)}$
- D.  $\text{Cl}_{2(g)}$ ,  $\text{H}_{2(g)}$ , and  $\text{NaOH}_{(aq)}$

**Use the following information to answer the next 3 questions**

The chlor-alkali process used by Dow Chemical in Fort Saskatchewan uses sodium chloride from underground deposits. The sodium chloride is dissolved in water and then pumped into electrolytic cells where a current is passed through the solution to form yellow chlorine gas, colorless hydrogen gas and aqueous sodium hydroxide. The ionic equation is



In the electrolysis of  $\text{NaCl}_{(aq)}$ , the cathode half-reaction is

- A.  $\text{Na}^+_{(aq)} + \text{e}^- \rightarrow \text{Na}_{(s)}$
- B.  $2 \text{Cl}^-_{(aq)} \rightarrow \text{Cl}_{2(g)} + 2 \text{e}^-$
- C.  $2 \text{H}_2\text{O}_{(l)} \rightarrow \text{O}_{2(g)} + 4 \text{H}^+_{(aq)} + 4 \text{e}^-$
- D.  $2 \text{H}_2\text{O}_{(l)} + 2 \text{e}^- \rightarrow \text{H}_{2(g)} + 2 \text{OH}^-_{(aq)}$

**Numerical Response**

If the mass of the element formed at the anode is 78.1 g, the mass of element formed at the cathode is \_\_\_\_\_ g.

(Record your answer to three digits on the answer sheet.)

When the electric current is switched off, the

- A. pH stops decreasing
- B. concentration of the  $\text{Na}^+_{(aq)}$  stops changing
- C. concentration of the  $\text{Cl}^-_{(aq)}$  stops decreasing
- D. concentration of the  $\text{H}_2\text{O}_{(l)}$  stops decreasing